

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Further Notes on the Snowy Owl in Ontario. — Since my letter of March 3, 1902, was published by Mr. Ruthven Deane, in his paper on the Snowý Owl, in the July 'Auk,' further information has made it evident that the migration in Ontario was a much more extensive one than I had at first imagined. During March the females disappeared and were replaced in April by the returning flight of light colored birds (males, as far as I was able to examine). A few remained about Toronto Marsh all through May, and a small light colored male was taken on June 7. It was in excellent condition and showed no trace of being a wounded bird.

Estimate of the number killed. — It soon became apparent that this migration was no ordinary flight as regards numbers, and as answers to my enquiries came in I saw that some other means of counting heads was necessary. I then had recourse to the number of artificial owl eyes used in Ontario during the migration. I was greatly helped by two facts; first, the almost total absence of Horned Owls from Ontario, or at least the territory affected by the migration, and was thus able to eliminate the possibility of many of the eyes being used for Horned Owls; secondly, nine-tenths of the eyes used by taxidermists, amateur or professional, in Ontario are bought from three firms in Toronto. In one case I went over all the orders and checked off the owl eyes; in another I got a careful estimate, and in the third I estimated the number from information as to the extra eyes imported to meet the demand. I found that not less than five hundred pairs of large owl eyes were sold in Ontario during this migration; and I believe the figure to be a low one, for not only were the regular sizes exhausted, but any yellow eye that could be made to do duty was used. From what I heard and saw I believe that less than half of the owls killed were mounted; and in going over the matter with Dr. Wm. Brodie I found that he too had concluded that one thousand was within the mark, though on different grounds .- J. H. Fleming, Toronto, Ontario.

An Addition to the Avifauna of Virginia.—In a collection of birds made during May, 1902, by the writer, in the Lake Drummond region of the Dismal Swamp, there is a specimen of Hairy Woodpecker which proves to be typical of the southern subspecies, *Dryobates villosus audubonii* (Swains.). Mr. William Palmer very kindly compared the specimen (& ad., taken May 22, 1902, Washington ditch, ½ mile northwest of Lake Drummond, Dismal Swamp, Nansemond County, Virginia; field number 32, coll. of J. W. D. Jr., Washington, D. C.) with material in the National Museum, and pronounces it referable to the southern race. Hitherto this form has not been taken further north than North Carolina. — John W. Daniel, Jr., Washington, D. C.

A new Foster-parent of the Cowbird.—On April 28, 1902, I found in an old log cabin a nest of Bewick's Wren, containing five fresh eggs of the owner and one fresh egg of the Cowbird. This species I do not find mentioned in any book, not even the late Maj. Chas. E. Bendire's monu-